

# Kangaroo Baby Massage: An Intervention which Improves the Perceived Maternal Self-Efficacy\*

\* Article derived from the doctoral thesis "Effects of the Kangaroo Baby Massage on the mother-child interaction at home", submitted to the Universidad Nacional de Colombia, 2023. Available at: <https://repositorio.unal.edu.co/handle/unal/84198> Funded Research. Margaret Mc Namara Education Grants. Research project: Effects of massage to the kangaroo infant in the home.

---

✉ **Nubia Castiblanco-López**

<https://orcid.org/0000-0003-1390-7923>  
Universidad Nacional de Colombia, Colombia  
ncastiblancol@unal.edu.co

**Fred Gustavo Manrique Abril**

<https://orcid.org/0000-0002-0723-3240>  
Universidad Nacional de Colombia, Colombia  
Universidad Pedagógica y Tecnológica de Colombia,  
Colombia  
fgmanriquea@unal.edu.co

**Lucy Marcela Vesga Gualdrón**

<https://orcid.org/0000-0003-1575-1782>  
Universidad Nacional de Colombia, Colombia  
lmvesgag@unal.edu.co

Received: 08/08/2023  
Sent to peers: 09/26/2023  
Approved by peers: 11/22/2023  
Accepted: 02/09/2024

**DOI: 10.5294/aqui.2024.24.2.2**

**Para citar este artículo / To reference this article / Para citar este artigo**

Castiblanco-López N, Manrique FG, Vesga LM. Kangaroo Baby Massage: An intervention which improves the perceived maternal self-efficacy. *Aquichan*. 2024;24(2):e2422. DOI: <https://doi.org/10.5294/aqui.2024.24.2.2>

**Theme:** Health care technologies

**Contribution to the discipline:** The Kangaroo Baby Massage (KBM) is a pivotal nursing practice originated in 1996 with the Mother Kangaroo Program at the San Pedro Claver Clinic and was subsequently refined through experiences in various countries (Senegal, Cuba, Vietnam, and England). This article derives from the doctoral dissertation titled: *Effects of Massage on the Kangaroo Baby on the mother-child Interaction at Home*, which characterizes the KBM and integrates it with Kathryn Barnard's theory. It represents a significant contribution to nursing knowledge and offers an effective intervention for early childhood and maternal health.

## Abstract

**Introduction:** The Kangaroo Baby Massage (KBM) is a technique designed for preterm and low birth weight newborns that eliminates the need for an incubator, providing a practical and beneficial option for mothers at home. **Objective:** To test the effect of the KBM on perceived maternal self-efficacy favoring the mother-infant interaction at home. **Materials and methods:** A pragmatic, double-blind, randomized clinical trial was conducted in three phases: 1) KBM integration with Kathryn Barnard's theory, 2) study design, and 3) implementation and outcome evaluation. Two groups were defined: The intervention group (KBM) and the control group (which used the kangaroo position without massage), each group consisted of 34 mother-infant pairs. Recruitment took place within an outpatient kangaroo program in Bogota, Colombia. Implementation and follow-up were conducted via teleconsultation, using the KBM video "Diary of My Kangaroo Baby" and simulators. The perceived maternal self-efficacy questionnaire was conducted at three points in time: prior to the study and on the 7<sup>th</sup> and 14<sup>th</sup> day after the study. The questionnaire was analyzed with a statistical design of longitudinal data in F<sub>1</sub>LDF<sub>1</sub> factorial experiments. **Results:** Homogeneous groups were defined based on sociodemographic variables and maternal-perinatal history. Mothers who applied the KBM technique showed higher scores in perceived maternal self-efficacy at both 7 and 14 days compared to the control group mothers [RTE (lower limit; upper limit) - day 7: control = 0.502 (0.437; 0.567) vs. KBM = 0.503 (0.426; 0.581), and day 14: control = 0.564 (0.482; 0.640) vs. KBM = 0.719 (0.650; 0.776)]. **Conclusion:** The KBM is a nursing intervention that enhances perceived maternal self-efficacy, fostering greater family involvement and strengthening mother-child interaction.

### Keywords (Source: DeCS)

Self-efficacy; massage; teleconsultation; kangaroo-mother method; mother-infant interaction; pragmatic clinical trial; nursing theory.

## 4 El masaje al bebé canguro, una intervención que mejora la autoeficacia materna percibida\*

\* Artículo derivado de tesis doctoral “Efecto del masaje al bebé canguro sobre la interacción madre-hijo en el hogar”, presentada a la Universidad Nacional de Colombia, 2023. Disponible en: <https://repositorio.unal.edu.co/handle/unal/84198> Investigación financiada. Margaret Mc Namara Education Grants. Proyecto de investigación: Efecto del masaje al bebé canguro en el hogar.

### Resumen

**Introducción:** el masaje al bebé canguro (MBC) es una técnica para recién nacidos pretérmino y bajo peso al nacer que no requiere incubadora y favorece a las madres en el hogar. **Objetivo:** probar el efecto del MBC sobre la autoeficacia materna percibida para favorecer la interacción madre-hijo en el hogar. **Materiales y métodos:** ensayo clínico aleatorizado pragmático, doble ciego, desarrollado en tres fases: 1) integración MBC y teoría de Kathryn Barnard, 2) diseño del estudio y 3) ejecución y resultados. Se definieron dos grupos: intervención (MBC) y control (posición canguro sin masaje), con 34 diadas madre-hijo en cada grupo. Reclutamiento realizado en un programa canguro ambulatorio de Bogotá, Colombia. La ejecución y el seguimiento fue hecha a través de teleconsulta, el video MBC “Diario de mi bebé canguro” y simuladores. Se aplicó el cuestionario de autoeficacia materna percibida en tres momentos: antes de iniciar el estudio y a los días 7 y 14 después del estudio. Se analizó con un diseño estadístico de datos longitudinales en experimentos factoriales F1LDF1. **Resultados:** fueron definidos grupos homogéneos en las variables sociodemográficas y de antecedentes materno-perinatales. Las madres que aplicaron el MBC registraron puntajes más altos de autoeficacia materna percibida a los 7 y 14 días, en comparación con las madres del control [RTE (límite inferior; límite superior) – día 7: control = 0.502 (0.437; 0.567) vs. MBC = 0.503 (0.426; 0.581), y día 14: control = 0.564 (0.482; 0.640) vs. MBC = 0.719 (0.650; 0.776)]. **Conclusión:** el MBC es una intervención de enfermería que mejoró la autoeficacia materna percibida, favoreciendo la participación de la familia y la interacción madre-hijo.

#### Palabras clave (Fuente: DeCS)

Autoeficacia; masaje; teleconsulta; método madre-canguro; interacción madre-hijo; ensayo clínico pragmático; teoría de enfermería.

# Massagem no bebê canguru, uma intervenção que melhora a autoeficácia materna percebida\*

Artigo derivado da tese de doutorado “Efecto del masaje al bebé canguro sobre la interacción madre-hijo en el hogar” (“Efeito da massagem no bebê canguru sobre a interação mãe-filho no domicílio”), defendida na Universidad Nacional de Colombia, 2023. Disponível em: <https://repositorio.unal.edu.co/handle/unal/84198> / Pesquisa financiada. Margaret Mc Namara Education Grants. Projeto de pesquisa “Efeito da massagem no bebê canguru no domicílio”.

## Resumo

**Introdução:** a massagem no bebê canguru (MBC) é uma técnica para bebês prematuros e de baixo peso que não necessita de incubadora e que dá apoio às mães no domicílio. **Objetivo:** testar o efeito da MBC na autoeficácia materna percebida para apoiar a interação mãe-bebê em casa. **Materiais e método:** ensaio clínico randomizado, pragmático, duplo-cego, desenvolvido em três fases: i) integração da MBC e da teoria de Kathryn Barnard, ii) desenho do estudo e iii) execução e resultados. Foram definidos dois grupos: intervenção (MBC) e controle (posição canguru sem massagem), com 34 díades mãe-bebê em cada grupo. O recrutamento foi realizado em um programa canguru ambulatorial em Bogotá, Colômbia. A implementação e o acompanhamento foram feitos por meio de teleconsultas, do vídeo MBC “Diário do meu bebê canguru” e de simuladores. O questionário de autoeficácia materna percebida foi aplicado em três momentos: antes do início do estudo e nos dias 7 e 14 após o estudo. Ele foi analisado com um projeto estatístico de dados longitudinais em experimentos fatoriais F1LDF1. **Resultados:** foram definidos grupos homogêneos para variáveis sociodemográficas e histórico materno-perinatal. As mães que aplicaram o MBC registraram pontuações mais altas de autoeficácia materna percebida aos 7 e 14 dias, em comparação com as mães do controle [RTE (limite inferior; limite superior) — dia 7: controle = 0,502 (0,437; 0,567) vs. MBC = 0,503 (0,426; 0,581), e dia 14: controle = 0,564 (0,482; 0,640) vs. MBC = 0,719 (0,650; 0,776)]. Conclusões: a MBC é uma intervenção de enfermagem que melhorou a autoeficácia materna percebida, favorecendo o envolvimento familiar e a interação entre mãe e filho.

### Palavras-chave (Fonte: DeCS)

Autoeficácia; massagem; teleconsulta; método mãe-canguru; interação mãe-filho; ensaio clínico pragmático; teoria de enfermagem.

## Introduction

The Kangaroo Mother Care (KMC) is an alternative to reduce stress for hospitalized babies and their mothers (1). However, several challenges can cause stress for mothers, such as fear of injuring their baby, difficulty maintaining the kangaroo position, fatigue, gender role concerns, limited opportunities for parents to practice the method (2), and lack of family support (3), negatively impacting maternal perceptions of their ability to care for their baby at home and affecting the quality of mother-child interaction.

Barnard's theory emphasizes that successful mother-child interaction relies on the ability of both to adapt to each other, with the quality of the socio-emotional environment in early childhood largely dependent on maternal competence in establishing an emotionally supportive and harmonious setting (4). The woman who can effectively meet her child's needs is considered a competent mother (5). However, preterm babies often provide limited responses to their mothers, and if the mother lacks sufficient understanding of her child's behavior, this dynamic can disrupt their interaction, leading her to perceive herself as incapable of alleviating her baby's discomfort (4).

The mother's ability to alleviate the baby's feelings of discomfort is one of the characteristics of mothers that Barnard argued for in her theory (4). Consequently, mothers who respond to and meet their children's needs gain confidence and competence, generating a positive interactive experience (4). However, when a premature baby does not send clear signals or is developmentally unresponsive (6), it can undermine the mother's ability to perform her parenting role safely and effectively, meaning the mother's confidence, competency, and satisfaction (7).

Low self-efficacy affects mothers' responses when they interact with their children (8), making it necessary to pay attention both to the mother's and the newborn's emotions (9). The perception of self-efficacy refers to an individual's judgment about their own abilities to perform tasks and handle situations with an expectation of success. It is not an inherent quality, but rather how one perceives their capacity to manage specific situations (10), understood as a dynamic and emergent attribute, subject to change depending on the demands of a particular task or situation. In this context, being a competent mother involves the ability to effectively respond to her children's needs. It requires a harmonious approach to foster an interaction that is productive and satisfying for both mother and child within a nurturing environment (5).

Therefore, if the environment for premature infants is made pleasant—resembling the safety and comfort of the womb—and mothers are actively involved in the development of care interventions, it can significantly improve the child's neurobehavioral outcomes in the long term. This serves as a motivating factor for mothers (11), especially when interventions are conducted in natural settings like

the home. A scoping review concluded that infant massage administered by mothers to their hospitalized premature babies decreased their stress, improving maternal-infant interaction, satisfaction, and maternal competence in the short term (12).

Several interventions aim to foster early interactions between parents and premature infants, including auditory, tactile, vestibular, and visual (ATVV) massage therapy (13). While this technique has been evaluated with hospitalized infants, its application at home for babies in the kangaroo position has not been explored extensively, which means maintaining the kangaroo position during such interventions without disruption, and in the case of Bogota, with an average temperature of around 13-14°C. Furthermore, some mothers with premature babies at home are hesitant to perform massages due to a lack of knowledge and express a need for guidance from nurses on suitable techniques for the home environment (14).

Educating mothers about stimulation techniques is an essential step toward enhancing person- and family-centered care (15), considering that interventions targeting premature and low birth weight newborns can have profound future impacts on child health (16). Moreover, employing multimodal nursing intervention strategies, supported by valid and reliable scales like the perceived maternal parental self-efficacy questionnaire adapted to the Colombian context, significantly benefits both maternal and child health (17).

Nursing is vital for effective KMC management (18) and its professionals become key guides for its implementation (19). The practice itself helps overcome fear and reinforces positive experiences (20) while encouraging self-care for mothers and their families. Most massage techniques in premature newborns were performed in incubators and neonatal units, evidencing an absence of studies that apply a massage technique for premature babies in the kangaroo position at home, under the care of their mothers and families (15, 21, 22).

Nevertheless, the KBM technique is an intervention that has been used by mothers since 1996, allowing them to care for their premature infants without the need for an incubator, both in neonatal units and home settings. A 2021 pre-experimental study involving 63 premature newborns from an outpatient kangaroo program in Bogota, Colombia, highlighted the safety of KBM on physiological parameters, demonstrating positive effects on heart rate, oxygen saturation, and temperature (23), although the study did not explore its impact on perceived maternal self-efficacy.

The objective of the present study was to investigate the effect of the KBM on a mother's ability to alleviate her child's discomfort, measured through perceived maternal self-efficacy, grounded in

the hypothesis that KBM, when applied by mothers, would positively influence their perceived ability to effectively comfort and care for their infants, thereby enhancing maternal self-efficacy.

## Methodology

This quantitative study employed a pragmatic randomized clinical trial design, evaluated using the PRECIS-2 tool. It was a parallel, double-blind trial involving two groups: An experimental group with mothers applying the KBM technique to their children, and a control group with mothers carrying their children in a kangaroo position without massage.

Mother-kangaroo baby pairs were randomly assigned to either group to prevent measurement bias. A biostatistician used computer-generated sequences through Stata to create a simple randomization scheme in blocks of size four. This resulted in 34 pairs in the KBM intervention group and 34 pairs in the kangaroo position control group without massage. To ensure blinding, sealed dark envelopes marked with codes from 1 to 68 were used, each corresponding to a consecutive assignment number. Inside each envelope was a slip indicating group assignment: A red slip for the KBM group or a blue slip for the control group. Additionally, the envelope contained information about the nurse responsible for conducting the teleconsultation. The opening of the envelopes was witnessed by the mother and an independent observer to maintain transparency.

Recruitment considered inclusion and exclusion criteria that allowed selection biases to be controlled. The inclusion criteria took into account that the baby was less than 37 weeks of gestational age or with low birth weight, cared for by its mother in the kangaroo position at home, with a gestational age greater than or equal to 30 weeks, age-corrected equal to or greater than 35 weeks at the time of recruitment, weight equal to or greater than 2100 g at the time of entry into the study, and controls in the Outpatient Kangaroo Mother Program of the Centro Policlínico del Olaya. Mothers with at least two days of training in said program, with social support, a family member's phone number, WhatsApp, and email.

Babies with a history of grade III or IV intraventricular hemorrhages, congenital malformations, or conditions requiring hospitalization during the study were excluded, as well as mothers who were hospitalized at the time of recruitment, re-hospitalized during the study, had cognitive problems or mental disorders, or had live twins.

The sample consisted of 34 mother-kangaroo baby pairs in each group, with a confidence level of 95 % and a significance level of 5 %. An estimated 10 % loss was accounted for. The independent variable was the KBM intervention, while the dependent variable was the total score of the perceived maternal self-efficacy questionnaire, which has robust psychometric properties for use with parents of neonatal populations. Initially validated with mothers of hospitalized premature newborns, the questionnaire has been



proven to be a reliable, robust, and valid psychometric measure (25). It has also been utilized in other studies during the postnatal period and with first-time mothers (26, 27).

Vargas et al. (26) were the first to translate this instrument into Spanish to use it in practice, research and, especially, easily measure maternal self-efficacy not only during the hospital stay but also at discharge and during postpartum follow-ups. The study found that the questionnaire had high face validity for women, with a comprehension rate of (99 %). For experts, it showed medium comprehension (84.1 %), medium clarity (83.9%), and medium precision (80 %). The content validity index was robust, with relevance scores of 0.93 and 0.96. The overall Cronbach's alpha value was 0.98 (95 % CI: 0.97-0.98). Therefore, the Spanish version of the questionnaire, with its robust properties, is suitable for identifying women with low self-efficacy and evaluating the effectiveness of nursing interventions.

The questionnaire contains twenty items, each rated on a Likert scale from 1 to 4 (strongly disagree, disagree, agree, strongly agree), with a maximum total score of 80. In this study, mothers completed the questionnaire through self-report at three points in time: before starting the intervention, 7 days after implementation, and 14 days after implementation. Additionally, there was a daily report noting who assisted with the kangaroo care and the mother's perception of "I am good at kangarooing" until the baby no longer required the position. These data were recorded in a printed manual titled *Diary of My Kangaroo Baby*, which also contained the perceived maternal self-efficacy questionnaires for days 7 and 14. Mothers were instructed to fill out the questionnaires and send photographic records via WhatsApp to the respective nurse conducting the follow-up. This method helped control measurement and information biases.

The information was collected in three stages:

1. Training of research assistants: A professional nurse expert in the Kangaroo Mother Program was trained virtually, who would be in charge of the control group; a research assistant was also trained in the management and systematization of the data, and a third research assistant was also trained to verify the data in its entirety.
2. Recruitment of the mother-baby kangaroo pairs: It was done in person, after the outpatient kangaroo program control. Once the mother voluntarily agreed to participate in the study and signed the informed and substituted consent, the sheets on sociodemographic aspects and maternal and perinatal history were completed, and the perceived maternal self-efficacy questionnaire was applied. The mother was trained in taking temperature with the baby in the kangaroo position and all mothers were given a kangaroo kit (the *Diary of My Kangaroo Baby* and black ink pen, 500 ml liquid hand

soap, cotton lycra kangaroo girdle, digital thermometer). Mothers assigned to the intervention group received the massage kit (a 100 ml bottle of sunflower oil and a soft styling brush).

3. Virtual Training and Home Implementation: The control group received training from a research assistant, who sent a connection link via WhatsApp. All mothers in this group were trained simultaneously at 8:00 p.m. through a nursing teleconsultation, which included a PDF presentation covering hand washing, correct use of face masks, kangaroo position, alarm signs, and management of the diary. Following the training, the mothers were instructed to take their baby's body temperature while in the kangaroo position, observe for ten minutes, take the temperature again, record it in the diary, and send a photo via WhatsApp. It is important to note that the control group did not receive any training related to massage and did not apply any type of massage. The nurse monitored the control group daily via WhatsApp until the baby no longer required the kangaroo position.

For the experimental group, the nurse sent mothers a connection link and an audiovisual guide titled "Kangaroo Baby Massage" via WhatsApp. Mothers were instructed to consult this guide and practice the KBM technique with a doll at home before the nursing teleconsultation. This audiovisual guide served as a step-by-step reference for mothers to apply the massage daily to their babies, available at <https://youtu.be/NV6ZDjodkXw?si=bf2fByrX6q6iwADI> (28).

The mother logged on at 8:00 p.m. from their home through their mobile phone or computer for the teleconsultation, a PDF presentation was used with the same topics as the control group; additionally, mothers in the intervention group received training on how to perform the KBM at home, they took the baby's temperature before and after giving the KBM, in a way that, while the researcher demonstrated through the screen the step by step of the application of the method on a simulator (doll), the mother was simultaneously performing it at home.

Mothers in both groups completed and reported the perceived maternal self-efficacy questionnaire at 7 and 14 days. The main researcher received the information after all the data were rigorously verified. Mother-child pairs were continuously followed and evaluated. The protocol included several safety measures, such as taking the temperature before and after the KBM intervention, and ensuring it was conducted in a secure room with closed windows and doors, without interrupting the kangaroo position. The mother always used the kangaroo girdle, with the baby remaining in position while the mother lay down with the head of the bed inclined at 30° to prevent the risk of falling. Additionally, all measures to minimize COVID-19 transmission were strictly followed, as synthesized in a literature review (29).

The study was classified as research with greater than minimal risk due to the therapeutic intervention; however, the KBM is a non-in-

vasive technique. Permission to use the perceived maternal parenting self-efficacy questionnaire was obtained, along with ethics approval from the Universidad Nacional de Colombia and the Centro Policlinico del Olaya. The study was registered prior to the pilot test on the United States National Library of Medicine platform (clinicaltrials.gov) with the identifier NCT04908332 (30).

## Results

The intervention was implemented and evaluated from August to December 2021 in the Outpatient Kangaroo Mother Program at the Centro Policlinico del Olaya in Bogota. At the time, the program was not conducted as a collective consultation due to the need to expand intensive care for COVID-19 patients and implement isolation measures. Consequently, the program operated in two offices where only the mother and her baby were allowed entry, adhering to protocols for mask use, hand washing, and social distancing. Meetings between mothers and group educational activities were strictly prohibited, minimizing the risk of sample contamination.


Figure 1 presents the flow chart according to the Consolidated Standards of Reporting Trials (CONSORT) (31), detailing the procedure for inclusion, assignment, monitoring, and analysis of the sample. The study was well-received and widely adhered to by participants.

The baseline data were homogeneous in the two groups (KBM vs. control), except for maternal age, finding that mothers in the KBM group were older ( $28 \pm 6.4$ ) than the mothers in the control group ( $24.4 \pm 5.1$ ) ( $p=0.014$ ). These data are presented in Tables 1, 2 and 3, corresponding to sociodemographic aspects, maternal history, and perinatal characteristics.

**Table 1.** Sociodemographic Aspects KBM vs. Control

Variable	KBM n=34 (%)	Control n=34 (%)	p
<b>Occupation</b>			0.806*
Housewife	20 (58.2)	19 (55.8)	
Employee, student	14	15	
<b>Monthly Income</b>			0.798*
Up to one minimum wage	22 (64.7)	23 (67.64)	
More than a minimum wage	12 (35.3)	11 (32.6)	
<b>Education Level</b>			0.948*
Up to incomplete high school	7	6	
Complete high school education	16	17	
University studies	11	11	
<b>Maternal Age (<math>\pm</math>SD) years</b>	28( $\pm$ 6.4)	24.4( $\pm$ 5.1)	0.014T
<b>Social Support Network</b>			1.00*
Partner, mother, or mother-in-law	27 (79.41)	27 (79.41)	
Other family members	7	7	
SD=Standard deviation *Chi square. T=t student			

Source: (32).



Source: (32).

**Table 2.** Maternal History KBM vs. Control

Variable	KBM n =34(%)	Control n =34(%)	p
<b>Desired pregnancy</b>			0.317F
Yes	19 (55.8)	23 (67.6)	
No	15 (44.2)	11 (32.4)	
<b>History of Kangaroo Children</b>			0.742F
0	29 (85.2)	28 (82.4)	
1	5 (14.8)	6 (17.6)	
<b>Type of Delivery</b>			0.584F
Vaginal birth without stitches	7 (20.6)	8 (23.6)	
Vaginal birth with stitches	8 (23.6)	5 (14.8)	
Caesarean section	13 (38.2)	11 (32.2)	
Caesarean section with Pomeroy	6 (17.6)	10 (29.4)	
<b>Days of Mother's Hospitalization in the Intensive Care Unit (ICU)</b>			0.061F
0	31 (91.2)	33 (97)	
2-7	3(8.8)	0	
8	0	1 (3)	

Variable	KBM n =34(%)	Control n =34(%)	p
<b>Mother Positive for COVID-19</b>			0.495F
No	25 (73.6)	24 (70.6)	
Recovered Suspected	9(26.4)	9 (26.4) 1 (3.0)	
<b>Number of Training Days in the Kangaroo Program</b>			
2 days	18(52.8)	24(70.6)	
3-13 days	16(47.2)	10(29.4)	
<b>First-Time Mothers</b>			0.624F
Yes	18 (52.9)	21 (61.8)	
No	16 (47.1)	13 (38.2)	
<b>Mothers Hospitalized in the ICU</b>			0.614F
Yes	3 (8,8)	1(3.0)	
No	31(91.2)	33 (97)	

Source: (32).

**Table 3.** Perinatal Characteristics KBM vs. Control

Variable	KBM	Control	Q
	N=34 (%)	N=34 (%)	
<b>Prematurity</b>			0.627F
Premature	19 (55.9)	17 (50)	
LBW Term	15 (44.1)	17 (50)	
Chronological age M ( ± SD)	<b>11.29 ( ± 7.53)</b>	<b>9.52 ( ± 6.39)</b>	0.307 U
<b>Corrected age</b>			0.159F
35-36.9	6 (17.6)	11(32.3)	
37-40	28 (82.4)	23 (67.6)	
<b>Sex</b>			0.791F
Male	9 (26.5)	11 (32.4)	
Female	25 (73.5)	23 (67.6)	
Birth weight in g M (±SD)	2179.55(±264.75)	2215.73(±207.7)	0.533T
<b>Adaptation type</b>			0.365F
Spontaneous	28 (82.4)	25(73.5)	
Driven Induced	3(8,8) 3(8,8)	7(20,6) 2(5,9)	
<b>Days of hospitalization in the NICU</b>			0.705F
0	28 (41.2)	29 (42.6)	
1-7	3(8,8)	3(8,8)	
8-14	2 (5.9)	2 (5.9)	
17	1 (3.0)	0	
<b>Oxygen dependent at home</b>			1,000*
No	28(82.4)	28(82.4)	
Yes	6 (17.6)	6 (17.6)	

Note: F=Fisher; \*Chi squared; U= Mann-Whitney; M (±SD) = Mean (± standard deviation); T= T student; NICU= Neonatal care unit.

Source: (32).

Table 4 presents the family support received by mothers in the KBM group, where a family member (typically the father or grandmother) took over kangarooing duties, allowing the mothers to rest. This support showed statistically significant differences over the days ( $p < 0.05$ ). In contrast, mothers in the control group generally performed kangaroo care alone. Table 5 shows that there were no statistically significant differences between the two groups ( $p > 0.05$ ) concerning the maternal perception expressed in the statement, "I am good at kangarooing my baby."

**Table 4 .** Family Support to Carry the Baby in the Kangaroo Position KBM vs. Control

Day	n	Group	Dad and Grandmother	Only the Mother	Other People	p
1	3.4	Control	1	11	22	0*
	3.4	KBM	20	1	13	
2	33	Control	1	9	23	0*
	33	KBM	18	1	13	
3	3.4	Control	4	7	23	0.01*
	33	KBM	16	3	14	
4	3.4	Control	1	9	24	0.001*
	31	KBM	15	2	14	
5	30	Control	0	7	23	0*
	30	KBM	15	4	11	
6	31	Control	2	8	21	0.01F
	28	KBM	14	3	11	
7	28	Control	2	7	19	0.001F
	24	KBM	14	3	7	
8	26	Control	2	7	17	0.024F
	20	KBM	9	3	8	
9	19	Control	1	6	12	0.002F
	18	KBM	11	2	5	
10	16	Control	1	5	10	0.033F
	15	KBM	7	1	7	

F = Fisher; \* Chi squared; KBM= Kangaroo Baby Massage; Significance 0.05

Source: (32).

**Table 5.** Maternal Perception: “I am good at kangarooing.” KBM vs. Control

Day	n	Group	Disagree	Agree	Strongly Agree	p
1	3.4	Control		14	20	1*
	3.4	KBM		14	20	
2	33	Control		17	16	0.215*
	33	KBM		12	21	
3	3.4	Control		17	17	0.056*
	33	KBM		9	24	
4	3.4	Control		15	19	0.206F
	31	KBM		9	22	
5	30	Control		11	19	0.584F
	30	KBM		9	21	
6	31	Control		13	18	0.283F
	28	KBM		8	20	
7	28	Control		12	16	0.115F
	24	KBM	1	8	15	
8	26	Control		8	18	0.187F
	twenty	KBM		3	17	
9	19	Control		7	12	0.162F
	18	KBM		3	15	
10	16	Control		5	11	0.256F
	fifteen	KBM	1	2	12	

Note: \*Chi = square; F = Fisher; KBM= Kangaroo Baby Massage (intervention): Significance 0.05

Source: (32).

The mother’s ability to alleviate the baby’s discomfort was assessed using the total score from the perceived maternal self-efficacy questionnaire, which was administered to 68 mothers with kangaroo babies. These mothers were divided into the intervention (KBM) and control groups, and the questionnaire was applied at three different points in time (days 0, 7, and 14). Table 6 indicates that, although mothers in the KBM group started with a lower score—median (Q1; Q3) = 68.5 (56; 80)—compared to mothers in the control group [71.5 (48; 79)], by day 14, the score for the KBM group increased considerably to 77.5 (66; 80), compared to 75 (61; 80) for the control group. Consequently, the final measurement of the questionnaire revealed a significant difference between the groups, with higher scores for the mothers who applied the KBM technique at home.

**Table 6.** Total Perceived Maternal Self-Efficacy score KBM vs. Control

Time	Group	n	Minimum	Maximum	Median	Q1	Q3	IQR	Half	D.S.
Day 0	KBM	3-4	56	80	68.5	64.25	72.75	8.5	68.176	5.697
Day 07	KBM	3-4	60	80	72.5	70	77	7	72.382	5.919
Day 14	KBM	3-4	66	80	77.5	75.25	80	4.75	76.735	3.752
Day 0	Control	3-4	48	79	71.5	64.25	75	10.75	69.324	8.112
Day 07	Control	3-4	55	80	72.5	68.25	78	9.75	72.324	6.153
Day 14	Control	3-4	61	80	75	68.25	79	10.75	73.5	6,121

Source: (32).

After running the F1LDF1 statistical model, Table 7 clearly shows that the relative treatment effects (RTE), with the respective 95 % confidence intervals, indicate a higher total score observed in the questionnaire. This change occurred significantly faster in the intervention group.

**Table 7.** RTE: Perceived Maternal Self-Efficacy in Time-Group Relationship KBM vs. Control)

Group	Time	n	Mean of the ranges	RTE	Bias	Variance	Lower limit	Upper limit
KBM	1(day 0)	34	64.029	0.311	-0.0005	0.083	0.250	0.386
KBM	2(day 07)	34	103.177	0.503	-0.0001	0.109	0.426	0.581
KBM	3(day 14)	34	147.279	0.719	0.0007	0.071	0.650	0.776
Control	1 (day 0)	34	82.118	0.400	-0.0007	0.088	0.333	0.473
Control	2(day 07)	34	102.941	0.502	-0.0002	0.076	0.437	0.567
Control	3 (day 14)	34	115.456	0.564	0.0009	0.114	0.482	0.640

Source: (32).



Figure 2 shows the significant difference between groups, displaying better results in the total score of the mothers who applied the KBM at time 3 (day 14).


In summary, the KBM technique applied by mothers had a favorable effect on the ability to alleviate the newborn's feeling of discomfort, expressed in perceived maternal parenting self-efficacy.

## Discussion

This study's findings show sufficient statistical evidence of the positive effect of the KBM on perceived maternal self-efficacy in mothers who applied the method compared to the control group of mothers who only kangarooed, with differences also found over time. The results were given in RTE (lower limit; upper limit), in the three measurements (days 0, 7 and 14), both for the KBM group [day 0: 0.311 (0.250; 0.386), day 7: 0.503 (0.426; 0.581), day 14: 0.719 (0.650; 0.776)], as well as for the control group [day 0: 0.400 (0.333; 0.473), day 7: 0.502 (0.437; 0.567), day 14: 0.564 (0.482; 0.640)].

Prematurity, compared to a full-term birth, generally had no effect on the perception of maternal self-efficacy (33). This finding supported the use of the self-efficacy questionnaire with mothers of premature newborns at home as well as allowing testing the positive effect of the KBM on maternal self-efficacy compared to the control group over time. The results of this study showed statistically significant differences between the two groups (KBM vs. control), with higher total scores at day 14 [mean ( $\pm$  SD): KBM = 76.7 ( $\pm$  3.7) vs. control = 73.5 ( $\pm$  6)], in alignment with the findings of Tristão (33), who reported no significant differences in mean total scores ( $\pm$  SD) between mothers of premature babies [65 ( $\pm$  9)] and mothers of full-term babies [65.5 ( $\pm$  8.2)]. Similarly, Vargas reported no significant differences in mean total scores ( $\pm$  SD) between first-time mothers of full-term babies receiving the multimodal intervention *Maternal support in the process of becoming a first-time mother* = 70.03 ( $\pm$  7.08) and those receiving usual postnatal care = 69.73 ( $\pm$  7.75);  $p=0.87$ . In a similar fashion, Taneja (34) found no significant effect on maternal competence scores at 6 weeks and 12 months after promoting skin-to-skin contact and breastfeeding through home visits (intervention group) compared to the control group. A recent study highlighted low perceived maternal self-efficacy in adolescent mothers due to the birth of a premature child, particularly when hospitalization was required (35).

This research is notable for being the first to evaluate mothers' perceptions of being "good at kangarooing," finding that the KBM intervention did not alter this perception at any point in time (Table 5). Perceived maternal self-efficacy and family support uniquely explained variations in parenting stress during the



Source: (32).

postpartum period, compounded by the challenges of the COVID-19 pandemic (36). The findings of this study, along with others (36, 37), indicate that the severe restrictions of the pandemic, which forced families to stay at home, made cohabitation with a partner beneficial for alleviating parenting stress.

Empowerment is crucial for competence in caregiving and facilitates the transition to home life, allowing parents to recognize their abilities and believe in themselves (38). Consequently, mothers who applied KBM felt more competent and perceived themselves as more empowered in caring for their children.

The study by Torres et al. found that 57.6 % of mothers with partner support reported high self-efficacy in breastfeeding (39). Similarly, in our study, mothers who applied the KBM technique received more help from their partners, who also undertook nursing duties, allowing the mothers to rest. As recommended by Rebolledo (8), it is crucial to address maternal tiredness or fatigue and implement interventions like KBM to enhance maternal well-being and mental health.

Additionally, in line with Kourouma et al. (3), who suggested designing strategies and interventions to improve the acceptance of KMC, our findings with the application of the KBM technique of-

fer an alternative solution to minimize barriers related to gender-related fatigue (40), lack of family support (41), resistance from grandmothers, and cultural aspects that hinder the adoption of KMC (42, 43).

## Conclusions

The KBM technique significantly enhanced mothers' responsiveness to their infants' discomfort, thereby improving perceived self-efficacy compared to control mothers. The study revealed statistically significant differences between the groups over time [RTE (lower limit; upper limit) - Day 7: Control = 0.502 (0.437; 0.567) vs. KBM = 0.503 (0.426; 0.581); Day 14: Control = 0.564 (0.482; 0.640) vs. KBM = 0.719 (0.650; 0.776)]. Mothers in the KBM group reported higher scores on the perceived maternal-parental self-efficacy questionnaire at 14 days compared to those in the control group. Additionally, KBM strengthened family support ( $p < 0.05$ ) without affecting perceptions of "being good at kangarooing." Thus, KBM serves as a nursing intervention that, when applied by mothers to their infants, effectively alleviates infant discomfort and enhances perceived maternal parental self-efficacy.

Mothers and families with preterm infants often experience fatigue and stress at home, and KBM emerges as an effective intervention to bolster maternal self-efficacy, thereby reducing barriers to adopting KMC. This study underscores the positive impact of the KBM technique, grounded in Barnard's theory, and underscores the importance of ongoing research to further enrich nursing practices.

**Acknowledgments:** My sincere thanks to all the mothers, families, and kangaroo babies, as well as the entire research team and healthcare personnel at the Centro Policlínico del Olaya and to the Universidad Nacional de Colombia.

**Limitations:** Medium-term follow-up was not feasible due to infants being discharged and referred to different institutions for Phase 2 of the Mother Kangaroo Program, which posed challenges in tracking and incurred substantial costs.

**Conflict of Interest Statement:** None declared.

- Cañadas DC, Carreño TP, Borja CS, Perales AB. Benefits of Kangaroo Mother Care on the Physiological Stress Parameters of Preterm Infants and Mothers in Neonatal Intensive Care. *Int J Environ Res Public Health*. 2022;19(12).
- Cattaneo A, Amani A, Charpak N, De Leon-Mendoza S, Moxon S, Nimbalkar S, et al. Report on an international workshop on kangaroo mother care: Lessons learned and a vision for the future. *BMC Pregnancy Childbirth*. 2018;18(1):1-10.
- Kourouma KR, Agbré-Yacé ML, Doukouré D, Cissé L, Some-Méazieu C, Ouattara J, et al. Barriers and facilitators to kangaroo mother care implementation in Cote d'Ivoire: a qualitative study. *BMC Health Serv Res*. 2021;21(1211):1-15.
- Oxford ML, Findlay DM. Development of the Feeding and Teaching Scales. 2nd ed. En *NCAST Caregiver/Parent-Child Interaction Feeding Manual*. Washington: University of Washington, NCAST Publication; 2015. 3-28 p.
- Garay-gordovil MZ. Autoeficacia materna percibida y actitud de soporte en la interacción madre-hijo. concepto, medición y relaciones entre sí. *Miscelánea Comillas Rev Ciencias Humanas y Soc*. 2013;71(139):419-44.
- Lugo-Gil J, Tamis-LeMonda CS. Family resources and parenting quality: Links to children's cognitive development across the first 3 years. *Child Dev*. 2008;79:1065-85. DOI: <https://doi.org/10.1111/j.1467-8624.2008.01176.x>
- Barraza C, Jofré V, Ramírez J. Percepciones asociadas al método madre canguero de madres con recién nacidos prematuros. *Cienc y Enferm*. 2020;26(32):1-10. DOI: <https://doi.org/10.29393/CE26-26PACB30026>
- Andrade Rebolledo D, Vicente Parada B. fatiga postparto: revisión de la literatura. *Rev Chil Obs Ginecol*. 2018;83(2):161-9.
- Matassini-Eyzaguirre SM, Cam-Chang L, Fernández-Sierra C. Barreras para la implementación del Método Mamá Canguero. *Rev del Cuerpo Médico Hosp Nac Almanzor Aguinaga Asenjo*. 2021;14(3):297-303. DOI: <https://doi.org/10.35434/rcmh-naaa.2021.143.1248>
- Bandura A. Regulation of Cognitive Processes Through Perceived Self-Efficacy. *Dev Psychol*. 1989;25(5):729-35. DOI: <https://doi.org/10.1037/0012-1649.25.5.729>
- Shattawi KK. Suspended liminality: breastfeeding and becoming a mother in two NICUs. *Int J Adv Nurs Stud*. 2015;4(2):75-84. DOI: <https://doi.org/10.14419/ijans.v4i2.4877>
- McCarty DB, Willett S, Kimmel M, Dusing SC. Benefits of maternally-administered infant massage for mothers of hospitalized preterm infants: a scoping review. *Matern Heal Neonatol Perinatol*. 2023;9(1):1-16. DOI: <https://doi.org/10.1186/s40748-023-00151-7>
- Fernandez X. Revisión sistemática de intervenciones tempranas en bebés prematuros para fomentar las interacciones sensibles padres-bebé y el vínculo de apego. *Clínica Contemp*. 2022;13(1):2003-5. DOI: <https://doi.org/10.5093/cc2022a3>
- Castiblanco López N, Muñoz de Rodríguez L. Vision de las madres en el cuidado del hijo prematuro en el hogar. *Av en Enferm*. 2011;29(1):120-9. <http://revistas.unal.edu.co/index.php/avenferm/article/view/35865>
- Hima J. Effectiveness of oral motor stimulation administered by mothers of preterm infants- A pilot study. *J Neonatal Nurs*. 2018;24(5):261-5. DOI: <https://doi.org/10.1016/j.jnn.2018.05.001>
- Silveira RC, Mendes EW, Fuentesfria RN, Valentini NC, Procianny RS. Early intervention program for very low birth weight preterm infants and their parents: A study protocol. *BMC Pediatr*. 2018;18(1):1-11. DOI: <https://doi.org/10.1186/s12887-018-1240-6>
- Vargas C, Díaz ZMR. Efficacy of a multimodal nursing intervention strategy in the process of becoming a mother: A randomized controlled trial. *Res Nurs*. 2021;1-14.
- Torres Castro YS. Rol de enfermería en el manejo eficaz del método canguero. Guayaquil: Universidad de Guayaquil; 2022. <http://repositorio.ug.edu.ec/handle/redug/63117>
- Díaz Jacanamijoi R, Romero Quiñones D. Método canguero aplicado por el profesional de enfermería en la unidad de cuidados intensivos pediátrica y neonatal. *Salud, Arte y Cuid*. 2020;13(1):51-56 <https://dialnet.unirioja.es/servlet/articulo?codigo=8633580>
- Malakouti J, Jabraeeli M, Valizadeh S, Babapour J. Mothers' experience of having a preterm infant in the Neonatal Intensive Care Unit, a Phenomenological Study. *Iran J Crit Care Nurs*. 2013;5(4):172-81. <https://pdfs.semanticscholar.org/ao70/3cb-2f49a557e120270a17207f7bc8b061b3c.pdf>
- Aldana AC, Tessier R, Charpak N, Tarabulsky G. Randomised controlled trial on the impact of kinesthetic stimulation on early somatic growth of preterm infants in Kangaroo position. *Acta pediátrica*. 2018;108(7):1-7. DOI: <https://doi.org/10.1111/apa.14675>
- dos Anjos FR, Nakato AM, Hemberger PK, Nohama P, Sarquis ALF. Effects of hydrotherapy and tactile-kinesthetic stimulation on weight gain of preterm infants admitted in the Neonatal Intensive Care Unit. *J Pediatr (Rio J)*. 2022;98(2):155-60. DOI: <https://doi.org/10.1016/j.jpmed.2021.04.011>
- Díaz Ramos L. Respuesta fisiológica con la técnica Masaje al Bebe Canguero, en un programa ambulatorio de Bogotá. Bogotá: Universidad Nacional de Colombia; 2022.
- Ciapponi A. Herramienta para evaluar cuán pragmático es un ensayo clínico : PRECIS-2. *Evid Actual en práctica ambulatoria*. 2015;17(4):114-7. DOI: <https://doi.org/10.51987/evidencia.v17i4.6317>
- Barnes CR, Adamson-Macedo EN. Perceived Maternal Parenting Self-Efficacy (PMP S-E) tool: Development and validation with mothers of hospitalized preterm neonates. *J Adv Nurs*. 2007;60(5):550-560. DOI: <https://doi.org/10.1111/j.1365-2648.2007.04445.x>
- Vargas-Porras C, Roa-Díaz ZM, Barnes C, Adamson-Macedo EN, Ferré-Grau C, De Molina-Fernández MI. Psychometric Properties of the Spanish Version of the Perceived Maternal Parenting Self-efficacy (PMP S-E) Tool for Primiparous Women. *Matern Child Health J*. 2020;24(5):537-45. DOI: <https://doi.org/10.1007/s10995-019-02860-y>
- Aliabadi F, Borimnejad L, Kamali M, Rassafiani M, Nazi S. Perceived maternal parenting self-efficacy: Translation and face validation with Iranian mothers of hospitalized preterm neonates. *Iran Rehabil J*. 2013;11(Specialissue):7-10.
- Castiblanco-López N. Masaje al Bebé Canguero [YouTube]. Bogotá: Universidad Nacional de Colombia; 2017. [https://www.youtube.com/watch?v=qbhGj1aPKGk&t=4s&ab\\_channel=NunaEducaUniversidadNacionalDeColombia](https://www.youtube.com/watch?v=qbhGj1aPKGk&t=4s&ab_channel=NunaEducaUniversidadNacionalDeColombia)
- Castiblanco N. Lactancia materna y Programa Madre Canguero una simbiosis necesaria en tiempos de covid-19. In: *Lactancia materna en tiempos de covid-19*. Cartagena: Red INLAMA; 2021. p. 77-88.
- Castiblanco-López N. Effect of Kangaroo Baby Massage on Mother-infant Interaction at Home (KBM); 2021. <https://classic.clinicaltrials.gov/ct2/show/NCT04908332>
- Cobos-Carbó A, Augustovski F. Declaración CONSORT 2010: actualización de la lista de comprobación para informar ensayos clínicos aleatorizados de grupos paralelos. *Medicina Clínica*. 2011;137(5):213-215. DOI: <https://doi.org/10.1016/j.medcli.2010.09.034>

32. Tristão RM, Neiva ER, Barnes CR, Adamson-Macedo E. Validation of the scale of perceived self-efficacy of maternal parenting in Brazilian sample. *J Hum Growth Dev.* 2015;25(3):277-86. DOI: <https://doi.org/10.7322/jhgd.96759>
33. Taneja S, Sinha B, Upadhyay RP, Mazumder S, Sommerfelt H, Martines J, et al. Community initiated kangaroo mother care and early child development in low birth weight infants in India—a randomized controlled trial. *BMC Pediatr.* 2020;20(150):1-12. DOI: <https://doi.org/10.1186/s12887-020-02046-4>
34. Ramírez Robles M, Herrera Paredes JM, Moreno González MM. Nivel de autoeficacia materna en adolescentes de la región nor-este de Guanajuato, México. *ACC CIETNA Rev la Esc Enfermería.* 2022;9(2):82-93. DOI: <https://doi.org/10.35383/cietna.v9i2.854>
35. Lin HC, Zehnah PL, Koire A, Mittal L, Erdei C, Liu CH. Maternal Self-Efficacy Buffers the Effects of COVID-19–Related Experiences on Postpartum Parenting Stress. *JOGNN - J Obstet Gynecol Neonatal Nurs.* 2022;51(2):177-94. DOI: <https://doi.org/10.1016/j.jogn.2021.12.004>
36. Miller JJ, Cooley ME, Mihalec-Adkins BP. Examining the Impact of COVID-19 on Parental Stress: A Study of Foster Parents. *Child Adolesc Soc Work J.* 2022;39(2):147-56. DOI: <https://doi.org/10.1007/s10560-020-00725-w>
37. Galeano SPO, Maya AMS. Empowering parents to care for preterm infants. *Rev Cuid.* 2022;13(2):1-19.
38. Torres IL, Silva K, Gomes da Silva K, Torreglosa M, Goulart B, Parreira BDM. Autoeficacia en la lactancia materna, síntomas de ansiedad y factores asociados. *Rev Familia, Ciclos Vida e Saúde no Context Soc.* 2021;9(3):642-50. [https://www.redalyc.org/journal/4979/497970304013/497970304013\\_1.pdf](https://www.redalyc.org/journal/4979/497970304013/497970304013_1.pdf)
39. Jamali QZ, Id RS, Shahid F, Fatima A, Khalsa S, Spacek J, et al. Barriers and enablers for practicing kangaroo mother care (KMC) in rural Sindh, Pakistan. *PLoS One.* 2019;14(6):1-15. DOI: <https://doi.org/10.1371/journal.pone.0213225>
40. Kinshella MW, Hiwa T, Pickerill K, Vidler M, Dube Q, Goldfarb D, et al. Barriers and facilitators of facility-based kangaroo mother care in sub-Saharan Africa: a systematic review. *Pregnancy and Child-birth.* 2021;3:1-10. DOI: <https://doi.org/10.1186/s12884-021-03646-3>
41. Yue J, Liu J, Williams S, Zhang B, Zhao Y, Zhang Q, et al. Barriers and facilitators of kangaroo mother care adoption in five Chinese hospitals: a qualitative study. *BMC Public Health.* 2020;20:1-11. DOI: <https://doi.org/10.1186/s12889-020-09337-6>
42. Ortiz Y, Rojas J. Cultural Care Practices Provided at Home by the Zenú Indigenous Mothers to their Premature Children and to Those with Low Birth Weight. *Invest Educ Enferm.* 2022;40(2):e09. DOI: <https://doi.org/10.17533/udea.iee.v40n2e09>
43. Ortiz Y RJ. Cultural Care Practices Provided at Home by the Zenú Indigenous Mothers to their Premature Children and to Those with Low Birth Weight. *Invest Educ Enferm.* 2022;40(2):e09. DOI: <https://doi.org/10.17533/udea.iee.v40n2e09>